

REVIEW

Open Access

Mechanics rules cell biology

James HC Wang*¹ and Bin Li^{2,3}

Abstract

Cells in the musculoskeletal system are subjected to various mechanical forces *in vivo*. Years of research have shown that these mechanical forces, including tension and compression, greatly influence various cellular functions such as gene expression, cell proliferation and differentiation, and secretion of matrix proteins. Cells also use mechanotransduction mechanisms to convert mechanical signals into a cascade of cellular and molecular events. This mini-review provides an overview of cell mechanobiology to highlight the notion that mechanics, mainly in the form of mechanical forces, dictates cell behaviors in terms of both cellular mechanobiological responses and mechanotransduction.

1. Introduction

Mechanical forces act on humans at different levels, from the body as a whole to individual organs, tissues, and cells. It is well known that appropriate mechanical loads are beneficial to bone and muscle by enhancing their mass and strength. On the other hand, excessive mechanical forces can also be detrimental; for example, excessive mechanical loading of tendons plays a major role in the development of tendinopathy [1,2]. Thus, mechanical forces have a profound effect on tissue homeostasis and pathophysiology. The central players in the human body's response to mechanical forces are various types of mechano-sensitive cells. Examples of such cells include tenocytes in tendons, fibroblasts in ligaments and skin, osteocytes in bone, chondrocytes in articular cartilage, and endothelial cells in blood vessels. Mechanical forces induce a wide range of cellular events, including proliferation, differentiation, and gene and protein expression by both adult differentiated and stem cells [3]. This mini-review provides a concise overview of cellular mechanobiological responses, with a focus on cells from musculoskeletal tissues. In addition, mechanotransduction mechanisms, by which cells "convert" mechanical forces into cellular biochemical events, are also briefly reviewed to emphasize the notion that mechanics, mainly in the form of external and internal mechanical forces, plays a vital role in cell biology. Note that readers who are inter-

ested in a more broad and in-depth understanding of the role of mechanics in cell biology should consult relevant papers, which are abundant in the literature.

2. External Mechanical Forces

External mechanical forces are defined as forces, such as tensile, compressive, or shear stresses, that are applied to cells from their environment. Depending on the cell type, the forces can come in one form or a combination of them. For example, fibroblasts in tendons and ligaments are mainly under tensile stress *in vivo*, while chondrocytes and osteocytes are subjected to compression and shear stress due to fluid flow in addition to tensile forces. In blood vessels, endothelial cells lining the vessel surface are subjected to a combination of tensile stress due to vessel expansion, hydrostatic pressure, and fluid shear stress.

Because of the ability to control experimental conditions, *in vitro* model systems have been developed to investigate cellular mechanobiological responses. In many of these systems, tensile forces are applied to the substrate and hence cause substrate deformation, which in turn loads cells that adhere to the underlying substrate. There are two ways to apply tensile mechanical forces to cells: the substrate may be stretched uniaxially or biaxially. Uniaxial stretching is appropriate for application of mechanical forces to cells originating from tendons (e.g., patellar and Achilles tendons) and ligaments (e.g., anterior cruciate ligament and medial collateral ligament), as these cells are aligned with their long axis parallel to the tendon or ligament and are therefore subjected primarily to uniaxial stretching *in vivo* [4-6]. On the other hand,

* Correspondence: wanghc@pitt.edu

¹ MechanoBiology Laboratory, Department of Orthopaedic Surgery, University of Pittsburgh School of Medicine, 210 Lothrop St, BST E1640, Pittsburgh, PA 15213, USA

Full list of author information is available at the end of the article

biaxial stretching is applied to cells that are subjected to tensile forces in all directions *in vivo*, e.g., dermal fibroblasts. A few biaxial stretching systems have been devised, which typically use circular elastic membranes to produce isotropic strains independent of stretching direction [7-10].

Besides tensile forces, compressive forces can also be applied to cells that are subjected to compression *in vivo*. One way of applying compressive forces is through application of hydrostatic pressure [11,12]. Another technique uses direct platen abutment to apply compressive forces to cells. This type of loading system includes unconfined compression, in which constant or low-cycle intermittent loads are delivered by manually applying weights [13,14], and confined compression, in which cells are placed between two platens that are contained by a confining chamber [15-17]. These compressive loading systems can be used to investigate mechanobiological responses of cells in tissues primarily subjected to compression *in vivo*, such as articular cartilage.

While 2-D systems such as those above have provided us with much insightful information regarding cellular mechanobiological responses, they are inherently limited in that they cannot model an *in vivo* tissue environment where cells are surrounded by extracellular matrix (ECM). Therefore, 3-D systems have also been developed that enable cells to reside in a more *in vivo*-like environment, which better preserves cell phenotype. A widely used 3-D system in the area of wound healing research is a cell-populated collagen gel (CPCG) [18,19]. Cells such as fibroblasts exert contraction forces on the surrounding collagen gel, thus remodeling it [20,21]. External mechanical forces can also be applied to CPCGs to study cellular mechanobiological responses [22,23]. Similarly, bioartificial tissues (BATs) were developed to embed tendon cells in collagen gels [24]. The phenotype of tendon cells in BATs is better preserved than in 2-D systems where cells are attached to 2-D substrates. This is generally true for other types of cells as well; e.g. chondrocytes in 3-D cultures retain their phenotype whereas they quickly lose it when cultured in 2-D systems such as a plastic dish [25].

3. Internal Mechanical Forces

Internal mechanical forces are the forces generated by cells themselves and are usually referred to as intracellular tension [26,27]. In non-muscle cells, intracellular tension is generated by cross-bridging of actomyosin, a process powered by ATP hydrolysis [28,29]. Such tensile forces are then transmitted to the ECM via focal adhesions [30], and the forces acting on ECM are called cell traction forces (CTFs). CTFs play a vital role in cell mechanobiology, as they function to direct ECM assembly [31], control cell shape [32-34], permit cell movement [35-39], and maintain cellular tensional homeostasis [40-42].

CTFs also deform the ECM network and cause stress and strain in the network, which in turn modulate cellular functions such as gene expression and protein secretion [43,44]. Therefore, CTFs are critical in many fundamental biological processes such as embryogenesis, angiogenesis, and wound healing [3].

In general, mechanobiological investigations rely on cell-substrate adhesion to transmit external mechanical forces to cells. This is because external mechanical forces acting on cells can alter the equilibrium state of internal forces, thereby affecting cellular mechanobiological responses [45].

In addition to mechanical forces, other mechanics parameters such as substrate stiffness also have a profound influence on cell behavior. A striking example is that substrate stiffness alone can direct specific differentiation of human mesenchymal stem cells (hMSCs); soft substrates (0.1-1 kPa) mimicking brain tissues are neurogenic, whereas stiffer substrates (8-17 kPa) mimicking muscle are myogenic. Finally, even stiffer substrates (25-40 kPa) resembling osteoid matrix can induce hMSCs to undergo osteogenic differentiation [46,47].

4. Cellular Mechanobiological Responses

Depending on the type of cell and loading conditions, application of mechanical forces to cells affects a spectrum of cellular functions, including cell proliferation, differentiation, gene expression and protein synthesis of ECM components, and production of cytokines and growth factors. For instance, in one study, human tendon fibroblasts were shown to increase their proliferation as well as gene expression and protein production of type I collagen in a stretch magnitude-dependent manner [48]. In another study, when repetitive stretching at a magnitude of 5% and a frequency of 1 Hz was applied to human tendon fibroblasts for one day, cell proliferation increased significantly. When the same conditions were applied for two days, however, cell proliferation was inhibited [49], indicating that stretching-induced proliferation of tendon fibroblasts also depends on stretching duration. Finally, in human periodontal ligament fibroblasts, a 10% cyclic equi-biaxial compression decreased type I collagen mRNA expression and reduced synthesis of fibronectin as well as the amount of total protein; however, the same level of cyclic stretching increased type I collagen mRNA levels and total protein levels [50]. These findings show that tensile and compressive forces with the same magnitude induce differential cellular mechanobiological responses.

In addition to cell proliferation and protein expression, mechanical forces can also induce the expression and production of inflammatory mediators, including COX-2, PGE₂, and LTB₄, in a stretching magnitude-dependent fashion [6,51]. In the presence of IL-1 β , a potent inflam-

matory mediator present in injured tissues, 4% cyclic uniaxial stretching decreased COX-2 and MMP-1 gene expression and PGE₂ production whose levels had been elevated by IL-1 β treatment; in contrast, cells under 8% stretching further increased the expression levels of these genes and PGE₂ production in addition to the effects of IL-1 β stimulation [52]. The findings of this study indicate that mechanical loading regulates cellular inflammatory responses in a loading magnitude-dependent manner. These findings suggest that when tissues such as tendons are injured, appropriate levels of exercise could be beneficial as it may reduce the inflammatory response. On the other hand, excessive loading of injured tendons, which may worsen tissue inflammation, could be detrimental. In chondrocytes, mechanical loading has also been found to regulate cellular inflammatory response via the NF- κ B signaling pathway [53].

While numerous studies have focused on studying mechanobiological responses of adult cells, efforts have been placed in recent years on investigating mechanobiological responses of stem cells. Accumulating evidence has shown that mechanical forces regulate proliferation and differentiation of stem cells. For instance, various mechanical loads applied to bovine bone marrow stem cells (BMSCs) induce differentiation of the stem cells into different cell lineages, including ligament cells, chondrocytes, myocardial and vascular cells [54-56]. In addition, small-magnitude stretching promotes osteogenic differentiation of hMSCs, whereas large-magnitude stretching induces tenogenic differentiation, as evidenced by up-regulation of genes specific for osteogenesis and tenogenesis, respectively [57]. Moreover, recent studies have demonstrated that cyclic uniaxial stretching not only enhances proliferation of rabbit tendon stem cells (TSCs), but also induces TSC differentiation into tenocyte and non-tenocyte lineages in a loading magnitude-dependent manner [58].

Corresponding to *in vitro* findings, mechanical forces *in vivo*, usually in the form of exercise, also induce various effects on tissues. For instance, exercise increases procollagen expression, collagen synthesis, and interstitial TGF- β concentration in humans [59]. In mice, moderate treadmill running induces the presence of myofibroblasts in tendons, suggesting that active remodeling takes place in response to applied loading on the tendon [60]. On the other hand, excessive mechanical forces acting on tendons *in vivo* cause degenerative changes in tendons (tendinopathy) [1,2].

Since mechanical loads are essential for the development, function, and repair of body components, mechanical conditioning is used in tissue engineering for proper development and functioning of tissue replacement constructs, especially for those bearing mechanical loads *in vivo* [61,62]. These constructs usually consist of deform-

able 3-D matrices seeded with cells and can be mechanically loaded using specially designed setups. For example, application of uniaxial stretching to 3-D collagen matrices populated with tendon fibroblasts resulted in expression of type I collagen and fibronectin similar to that of native tendons. This indicates that tendon cells residing in a mechanically loaded 3-D construct could be assuming a similar phenotype as those cells in native tendons. Moreover, these constructs were mechanically stronger than their unloaded counterparts [24]. When autogenous tissue engineered constructs of the patellar tendon made of type I collagen sponges and rabbit MSCs were mechanically stimulated, the stiffness of the cell-collagen constructs increased by 1.5 times compared to unloaded constructs [63]. Also, in smooth muscle cell (SMC)-seeded scaffolds, cyclic mechanical stretching for 5-20 weeks stimulated gene expression of SMC elastin and type I collagen. Tensile strength and Young's moduli of constructs were increased by cyclic stretching for 20 weeks, whereas both decreased over time without mechanical stretching [64].

In addition to external mechanical forces, internal mechanical forces can also regulate cell biology in terms of anabolic or catabolic states. For example, reducing internal mechanical forces (or intracellular tension) by releasing collagen gels from attachment to an underlying substrate or using cytochalasin D, a drug that disrupts actin cytoskeleton, changes rat tendon cells from an anabolic to a catabolic state, as measured by the gene expression levels of type I collagen and interstitial collagenase, respectively [65]. Moreover, cells can also use their internal contractile forces to regulate their own proliferation and differentiation [66,67].

Theoretical modeling predicted that accumulation of mechanical stress happens in a layer of cells that adhere to each other, and cells may use the mechanical stress as a feedback signal for their division [67,68]. Using a micro-patterned cell aggregate model, concentrated internal mechanical stresses around the perimeter of a cell aggregate were shown to cause endothelial or epithelial cells to proliferate at the perimeter but not in its inner region, where lower mechanical stresses were present [69]. Furthermore, collective internal mechanical stresses were also found to regulate differentiation of a cell aggregate, resulting in spatial patterning of differentiated cells [70,71]. Finally, decreasing intracellular tension or internal mechanical force by limiting cell spreading area has been shown to result in apoptosis [72].

5. Cellular Mechanotransduction

In order for a cell to respond to mechanical forces, the mechanical forces must be converted into chemical signals inside the cell to elicit a cascade of cellular and

molecular events. Such a process is termed cellular mechanotransduction (Fig. 1).

While the mechanisms of cellular mechanotransduction are still not completely understood, it is generally accepted that external mechanical forces acting on ECM have to be transmitted into a cell through integrin-mediated adhesions [73,74]. Integrins, which contain both a large ECM domain responsible for binding substrates and a cytoplasmic domain, are the main adhesive receptors and mechanotransducers that link the cytoskeleton to the ECM [26,75]. Therefore, the ECM-integrin-cytoskeleton pathway plays a major role in the mechano-signaling process. In a "tensegrity" model, mechanical forces applied to the cell membrane are directly and immediately transmitted to the nucleus through the inter-connected cytoskeleton composed of actin filaments, microtubules, and intermediate filaments [76]. Such a model is supported by the finding that application of mechanical stress to integrins altered the cytoskeleton and activated gene expression in a stress-dependent manner [77-79]. Using a FRET-based cytosolic Src reporter in a living cell, local stress was shown to induce rapid activation (< 0.3 sec) of Src at remote cytoplasmic sites; thus, a pre-stressed cytoskeleton can rapidly transduce mechanical signals [80].

In addition to integrins and the cytoskeleton, G proteins also function as mechanotransduction molecules [81,82]. Another important component of cellular mechanotransduction is intracellular Ca^{2+} [83]. Mechanical stretching of fibroblasts and many other types of cells increases the levels of intracellular Ca^{2+} , which serves as a secondary messenger [84,85]. In addition, cellular mechanotransduction also involves stretch-activated ion channels (SACs) [86,87]. In response to applied mechanical stresses, SACs open to allow ions like Ca^{2+} , Na^+ , and K^+ to pass through, thus transducing mechanical signals into activation of intracellular signaling molecules [88]. Finally, recent studies have shown that primary cilia also play an important role in cellular mechanotransduction. In bone cells, for example, primary cilia translate fluid flow into cellular responses independent of SACs [89].

In addition to the roles of many cellular components such as integrin and cytoskeleton in cellular mechanotransduction, researchers are also beginning to understand the mechanisms of how mechanical forces are initially sensed by the cell. In adherent cells, force transmission is primarily dependent on the attachment of cells to ECM molecules such as collagen or fibronectin [90]. Therefore, ECM proteins may function as "force sensors." Mechanical stresses acting on ECM may unfold a domain




Figure 1 Schematic illustration of the "mechanical nature" of cellular mechanotransduction mechanisms. Mechanical forces (MF) can induce mechanotransduction by directly altering conformation of an extracellular matrix (ECM) protein and integrin configuration and transmitting forces to the cytoskeleton and nucleus, thus eventually affecting transcription and translation. Also, mechanical forces can unfold a domain of the extracellular protein (M) and expose a cryptic site that may serve as an activating ligand for a cell surface receptor, resulting in a series of signaling events. Also, when mechanical forces are applied to "force receptors" (FR), such as integrins and G proteins, they initiate signal transduction, resulting in transcription followed by translation. As a result, soluble factors are secreted into the ECM, which act on the receptor (R) and then initiate a cascade of signaling events. Note that double arrows indicate intracellular tensions in the actin filaments. (Modified with permission from Wang and Thampatty, **Fig. four** in *Encyclopedia of Biomaterials and Biomedical Engineering*, 2008, p.1783-1793, Taylor & Francis).

of the ECM protein, resulting in exposure of its cryptic site, which may serve as an activating ligand for an adjacent receptor [83]. This potential force-sensing mechanism is supported by the finding that small and large forces unfold the weakest domain and the most stable domain of fibronectin, respectively [91]. Besides the conformation change in an ECM protein due to applied external mechanical forces, the cytoskeletal force, or the internal mechanical force, controls $\alpha_5\beta_1$ integrin switching between relaxed and tensioned states. Such a switch directly controls the strength of $\alpha_5\beta_1$ -fibronectin bond by engaging the synergy site in fibronectin [92].

6. Conclusion

Mechanical forces are ubiquitous and are known to greatly influence physiology and pathophysiology in humans. Mechano-responsive cells are responsible for these mechano-effects, as years of intensive mechanobiology research have shown that external mechanical forces influence a wide spectrum of cellular events, including alterations in cell proliferation, differentiation, gene expression, and protein production. It is also now appreciated that internal mechanical forces generated by cells themselves regulate cell biology in terms of metabolic state, cell proliferation and differentiation, etc. Particularly, CTFs, which are the internal mechanical forces transmitted to ECM, regulate many vital cellular functions such as migration and ECM assembly.

The keys to understanding mechanical force-regulated cell biology are cellular mechanotransduction mechanisms by which cells "convert" mechanical force signals into biochemical signals in cells. The role of ECM proteins, integrins, and cytoskeleton in cellular mechanotransduction is now firmly established. Recent studies also point to predominant role of primary cilia in mechanical signal transduction. They also show that mechanical forces may cause mechanotransduction events by altering conformation of signaling molecules, thus affecting their activity and consequently eliciting a cascade of biochemical events such as gene expression.

The fact that mechanics plays a dominant role in cell biology provides a solid foundation and rationale for use of mechanics to improve human health by designing appropriate equipment/instruments, exercise protocols, and rehabilitation regimens. For instance, in sports medicine, such practices will help improve overall performance while reducing and preventing musculoskeletal injuries in athletes. Also, combined use of "bio-interventions" and "mechanics" will further improve the outcome of clinical treatments of musculoskeletal injuries.

Competing interests

The authors declare that they have no competing interests.

Authors' contributions

JW and BL drafted and revised the manuscript together. Both authors read and approved the final manuscript.

Acknowledgements

This work is supported in part by NIH AR049921, AR049921S1, and AR049921S2 (JHW).

Author Details

¹MechanoBiology Laboratory, Department of Orthopaedic Surgery, University of Pittsburgh School of Medicine, 210 Lothrop St, BST E1640, Pittsburgh, PA 15213, USA, ²Orthopedic Institute, Soochow University, 708 Renmin Rd, Suzhou, Jiangsu 215007, China and ³Department of Orthopedics, The First Affiliated Hospital of Soochow University, 188 Shizi St, Suzhou, Jiangsu 215006, China

Received: 27 March 2010 Accepted: 8 July 2010

Published: 8 July 2010

References

1. Fredberg U, Stengaard-Pedersen K: **Chronic tendinopathy tissue pathology, pain mechanisms, and etiology with a special focus on inflammation.** *Scand J Med Sci Sports* 2008, **18**(1):3-15.
2. Wang JH, Iosifidis MI, Fu FH: **Biomechanical basis for tendinopathy.** *Clin Orthop Relat Res* 2006, **443**:320-332.
3. Wang JH, Thampatty BP: **Mechanobiology of adult and stem cells.** *Int Rev Cell Mol Biol* 2008, **271**:301-346.
4. Archambault J, Tsuzaki M, Herzog W, Banes AJ: **Stretch and interleukin-1beta induce matrix metalloproteinases in rabbit tendon cells in vitro.** *Journal of Orthopaedic Research* 2002, **20**(1):36-39.
5. Kim SG, Akaike T, Sasagawa T, Atomi Y, Kurosawa H: **Gene expression of type I and type III collagen by mechanical stretch in anterior cruciate ligament cells.** *Cell Structure & Function* 2002, **27**(3):139-144.
6. Li Z, Yang G, Khan M, Stone D, Woo SL, Wang JH: **Inflammatory response of human tendon fibroblasts to cyclic mechanical stretching.** *American Journal of Sports Medicine* 2004, **32**(2):435-440.
7. Hung CT, Williams JL: **A Method for Inducing Equi-Biaxial and Uniform Strains in Elastomeric Membranes Used as Cell Substrates.** *Journal of Biomechanics* 1994, **27**(2):227-232.
8. Lee AA, Delhaas T, Waldman LK, MacKenna DA, Villarreal FJ, McCulloch AD: **An equibiaxial strain system for cultured cells.** *Am J Physiol* 1996, **271**(4 Pt 1):C1400-1408.
9. Schaffer JL, Rizen M, Litalien GJ, Benbrahim A, Megerman J, Gerstenfeld LC, Gray ML: **Device for the Application of a Dynamic Biaxially Uniform and Isotropic Strain to a Flexible Cell-Culture Membrane.** *Journal of Orthopaedic Research* 1994, **12**(5):709-719.
10. Sotoudeh M, Jalali S, Usami S, Shyy JY, Chien S: **A strain device imposing dynamic and uniform equi-biaxial strain to cultured cells.** *Ann Biomed Eng* 1998, **26**(2):181-189.
11. Angele P, Yoo JU, Smith C, Mansour J, Jepsen KJ, Nerlich M, Johnstone B: **Cyclic hydrostatic pressure enhances the chondrogenic phenotype of human mesenchymal progenitor cells differentiated in vitro.** *J Orthop Res* 2003, **21**(3):451-457.
12. Toyoda T, Seedhom BB, Yao JQ, Kirkham J, Brookes S, Bonass WA: **Hydrostatic pressure modulates proteoglycan metabolism in chondrocytes seeded in agarose.** *Arthritis & Rheumatism* 2003, **48**(10):2865-2872.
13. Burton-Wurster N, Vernier-Singer M, Farquhar T, Lust G: **Effect of compressive loading and unloading on the synthesis of total protein, proteoglycan, and fibronectin by canine cartilage explants.** *Journal of Orthopaedic Research* 1993, **11**(5):717-729.
14. Steinmeyer J, Torzilli PA, Burton-Wurster N, Lust G: **A new pressure chamber to study the biosynthetic response of articular cartilage to mechanical loading.** *Res Exp Med (Berl)* 1993, **193**(3):137-142.
15. Chen AC, Sah RL: **Effect of static compression on proteoglycan biosynthesis by chondrocytes transplanted to articular cartilage in vitro.** *J Orthop Res* 1998, **16**(5):542-550.
16. Davisson T, Kunig S, Chen A, Sah R, Ratcliffe A: **Static and dynamic compression modulate matrix metabolism in tissue engineered cartilage.** *Journal of Orthopaedic Research* 2002, **20**(4):842-848.

17. Freeman PM, Natarajan RN, Kimura JH, Andriacchi TP: **Chondrocyte cells respond mechanically to compressive loads.** *Journal of Orthopaedic Research* 1994, **12**(3):311-320.
18. Bell E, Ehrlich HP, Buttle DJ, Nakatsuji T: **Living tissue formed in vitro and accepted as skin-equivalent tissue of full thickness.** *Science* 1981, **211**(4486):1052-1054.
19. Bell E, Ivarsson B, Merrill C: **Production of a tissue-like structure by contraction of collagen lattices by human fibroblasts of different proliferative potential in vitro.** *Proc Natl Acad Sci USA* 1979, **76**(3):1274-1278.
20. Grinnell F: **Fibroblast-collagen-matrix contraction: growth-factor signalling and mechanical loading.** *Trends in Cell Biology* 2000, **10**(9):362-365.
21. Fernandez P, Bausch AR: **The compaction of gels by cells: a case of collective mechanical activity.** *Integr Biol (Camb)* 2009, **1**(3):252-259.
22. Akhouayri O, Lafage-Proust MH, Rattner A, Laroche N, Caillot-Augusseau A, Alexandre C, Vico L: **Effects of static or dynamic mechanical stresses on osteoblast phenotype expression in three-dimensional contractile collagen gels.** *J Cell Biochem* 1999, **76**(2):217-230.
23. Peperzak KA, Gilbert TW, Wang JH: **A multi-station dynamic-culture force monitor system to study cell mechanobiology.** *Med Eng Phys* 2004, **26**(4):355-358.
24. Garvin J, Qi J, Maloney M, Banes AJ: **Novel system for engineering bioartificial tendons and application of mechanical load.** *Tissue Engineering* 2003, **9**(5):967-979.
25. Yates KE, Allemann F, Glowacki J: **Phenotypic analysis of bovine chondrocytes cultured in 3D collagen sponges: effect of serum substitutes.** *Cell Tissue Bank* 2005, **6**(1):45-54.
26. Ingber D: **Integrins as mechanochemical transducers.** *Curr Opin Cell Biol* 1991, **3**(5):841-848.
27. Burridge K, Chrzanoska-Wodnicka M: **Focal adhesions, contractility, and signaling.** *Annu Rev Cell Dev Biol* 1996, **12**:463-518.
28. Kolega J, Janson LW, Taylor DL: **The role of solation-contraction coupling in regulating stress fiber dynamics in nonmuscle cells.** *J Cell Biol* 1991, **114**(5):993-1003.
29. Sanger JW, Sanger JM, Jockusch BM: **Differences in the stress fibers between fibroblasts and epithelial cells.** *J Cell Biol* 1983, **96**(4):961-969.
30. Balaban NQ, Schwarz US, Riveline D, Goichberg P, Tzur G, Sabanay I, Mahalu D, Safran S, Bershadsky A, Addadi L, et al.: **Force and focal adhesion assembly: a close relationship studied using elastic micropatterned substrates.** *Nat Cell Biol* 2001, **3**(5):466-472.
31. Lemmon CA, Chen CS, Romer LH: **Cell traction forces direct fibronectin matrix assembly.** *Biophys J* 2009, **96**(2):729-738.
32. Li F, Li B, Wang QM, Wang JH: **Cell shape regulates collagen type I expression in human tendon fibroblasts.** *Cell Motil Cytoskeleton* 2008, **65**(4):332-341.
33. Wang HB, Dembo M, Wang YL: **Substrate flexibility regulates growth and apoptosis of normal but not transformed cells.** *Am J Physiol Cell Physiol* 2000, **279**(5):C1345-1350.
34. Kumar S, Maxwell IZ, Heisterkamp A, Polte TR, Lele TP, Salanga M, Mazur E, Ingber DE: **Viscoelastic retraction of single living stress fibers and its impact on cell shape, cytoskeletal organization, and extracellular matrix mechanics.** *Biophys J* 2006, **90**(10):3762-3773.
35. Ananthkrishnan R, Ehrlicher A: **The forces behind cell movement.** *Int J Biol Sci* 2007, **3**(5):303-317.
36. Rosel D, Brabek J, Tolde O, Mierke CT, Zitterbart DP, Raupach C, Bicanova K, Kollmannsberger P, Pankova D, Vesely P, et al.: **Up-regulation of Rho/ROCK signaling in sarcoma cells drives invasion and increased generation of protrusive forces.** *Mol Cancer Res* 2008, **6**(9):1410-1420.
37. Beningo KA, Dembo M, Kaverina I, Small JV, Wang YL: **Nascent focal adhesions are responsible for the generation of strong propulsive forces in migrating fibroblasts.** *J Cell Biol* 2001, **153**(4):881-888.
38. Ingber DE: **Mechanobiology and diseases of mechanotransduction.** *Annals of Medicine* 2003, **35**(8):564-577.
39. Lee J, Leonard M, Oliver T, Ishihara A, Jacobson K: **Traction forces generated by locomoting keratocytes.** *J Cell Biol* 1994, **127**(6 Pt 2):1957-1964.
40. Sawhney RK, Howard J: **Molecular dissection of the fibroblast-traction machinery.** *Cell Motil Cytoskeleton* 2004, **58**(3):175-185.
41. Eckes B, Krieg T: **Regulation of connective tissue homeostasis in the skin by mechanical forces.** *Clin Exp Rheumatol* 2004, **22**(3 Suppl 33):S73-76.
42. Harris AK: **Cell motility and the problem of anatomical homeostasis.** *J Cell Sci Suppl* 1987, **8**:121-140.
43. Tranquillo RT, Durrani MA, Moon AG: **Tissue Engineering Science - Consequences of Cell Traction Force.** *Cytotechnology* 1992, **10**(3):225-250.
44. Harris AK, Stopak D, Wild P: **Fibroblast traction as a mechanism for collagen morphogenesis.** *Nature* 1981, **290**(5803):249-251.
45. Ingber DE: **Tensegrity: the architectural basis of cellular mechanotransduction.** *Annu Rev Physiol* 1997, **59**:575-599.
46. Engler AJ, Griffin MA, Sen S, Bonnemann CG, Sweeney HL, Discher DE: **Myotubes differentiate optimally on substrates with tissue-like stiffness: pathological implications for soft or stiff microenvironments.** *J Cell Biol* 2004, **166**(6):877-887.
47. Rehfeldt F, Engler AJ, Eckhardt A, Ahmed F, Discher DE: **Cell responses to the mechanochemical microenvironment—implications for regenerative medicine and drug delivery.** *Adv Drug Deliv Rev* 2007, **59**(13):1329-1339.
48. Yang GG, Crawford RC, Wang JHC: **Proliferation and collagen production of human patellar tendon fibroblasts in response to cyclic uniaxial stretching in serum-free conditions.** *Journal of Biomechanics* 2004, **37**(10):1543-1550.
49. Barkhausen T, van Griensven M, Zeichen J, Bosch U: **Modulation of cell functions of human tendon fibroblasts by different repetitive cyclic mechanical stress patterns.** *Exp Toxicol Pathol* 2003, **55**(2-3):153-158.
50. He Y, Macarak EJ, Korostoff JM, Howard PS: **Compression and tension: differential effects on matrix accumulation by periodontal ligament fibroblasts in vitro.** *Connect Tissue Res* 2004, **45**(1):28-39.
51. Wang JH, Jia F, Yang G, Yang S, Campbell BH, Stone D, Woo SL: **Cyclic mechanical stretching of human tendon fibroblasts increases the production of prostaglandin E2 and levels of cyclooxygenase expression: a novel in vitro model study.** *Connective Tissue Research* 2003, **44**(3-4):128-133.
52. Yang G, Im HJ, Wang JH: **Repetitive mechanical stretching modulates IL-1 beta induced COX-2, MMP-1 expression, and PGE2 production in human patellar tendon fibroblasts.** *Gene* 2005, **363**:166-172.
53. Agarwal S, Deschner J, Long P, Verma A, Hofman C, Evans CH, Piesco N: **Role of NF-kappaB transcription factors in antiinflammatory and proinflammatory actions of mechanical signals.** *Arthritis Rheum* 2004, **50**(11):3541-3548.
54. Altman GH, Horan RL, Martin I, Farhadi J, Stark PR, Volloch V, Richmond JC, Vunjak-Novakovic G, Kaplan DL: **Cell differentiation by mechanical stress.** *FASEB J* 2002, **16**(2):270-272.
55. Park JS, Huang NF, Kurpinski KT, Patel S, Hsu S, Li S: **Mechanobiology of mesenchymal stem cells and their use in cardiovascular repair.** *Front Biosci* 2007, **12**:5098-5116.
56. Huang H, Kamm RD, Lee RT: **Cell mechanics and mechanotransduction: pathways, probes, and physiology.** *Am J Physiol Cell Physiol* 2004, **287**(1):C1-11.
57. Chen YJ, Huang CH, Lee IC, Lee YT, Chen MH, Young TH: **Effects of cyclic mechanical stretching on the mRNA expression of tendon/ligament-related and osteoblast-specific genes in human mesenchymal stem cells.** *Connect Tissue Res* 2008, **49**(1):7-14.
58. Zhang J, Wang JH: **Mechanobiological response of tendon stem cells: Implications of tendon homeostasis and pathogenesis of tendinopathy.** *J Orthop Res* 2010, **28**(5):639-643.
59. Kjaer M, Langberg H, Heinemeier K, Bayer ML, Hansen M, Holm L, Doessing S, Kongsgaard M, Krogsgaard MR, Magnusson SP: **From mechanical loading to collagen synthesis, structural changes and function in human tendon.** *Scand J Med Sci Sports* 2009, **19**(4):500-510.
60. Szczodry M, Zhang J, Lim C, Davitt HL, Yeager T, Fu FH, Wang JH: **Treadmill running exercise results in the presence of numerous myofibroblasts in mouse patellar tendons.** *J Orthop Res* 2009, **27**(10):1373-1378.
61. Butler DL, Juncosa-Melvin N, Boivin GP, Galloway MT, Shearn JT, Gooch C, Awad H: **Functional tissue engineering for tendon repair: A multidisciplinary strategy using mesenchymal stem cells, bioscaffolds, and mechanical stimulation.** *J Orthop Res* 2008, **26**(1):1-9.
62. Guilak F, Butler DL, Goldstein SA: **Functional tissue engineering: the role of biomechanics in articular cartilage repair.** *Clinical Orthopaedics & Related Research* 2001:S295-305.
63. Juncosa-Melvin N, Shearn JT, Boivin GP, Gooch C, Galloway MT, West JR, Nirmalanandhan VS, Bradica G, Butler DL: **Effects of mechanical stimulation on the biomechanics and histology of stem cell-collagen**

- sponge constructs for rabbit patellar tendon repair. *Tissue Eng* 2006, **12**(8):2291-2300.
64. Kim BS, Nikolovski J, Bonadio J, Smiley E, Mooney DJ: **Engineered smooth muscle tissues: regulating cell phenotype with the scaffold.** *Exp Cell Res* 1999, **251**(2):318-328.
65. Lavagnino M, Arnoczky SP: **In vitro alterations in cytoskeletal tensional homeostasis control gene expression in tendon cells.** *Journal of Orthopaedic Research* 2005, **23**(5):1211-1218.
66. Ingber DE: **Mechanical control of tissue morphogenesis during embryological development.** *Int J Dev Biol* 2006, **50**(2-3):255-266.
67. Shraiman BI: **Mechanical feedback as a possible regulator of tissue growth.** *Proc Natl Acad Sci USA* 2005, **102**(9):3318-3323.
68. Day C: **Mechanical force may determine the final size of tissues.** *Phys Today* 2007, **60**(4):20-21.
69. Nelson CM, Jean RP, Tan JL, Liu WF, Sniadecki NJ, Spector AA, Chen CS: **Emergent patterns of growth controlled by multicellular form and mechanics.** *Proc Natl Acad Sci USA* 2005, **102**(33):11594-11599.
70. Li B, Li F, Puskar KM, Wang JH: **Spatial patterning of cell proliferation and differentiation depends on mechanical stress magnitude.** *J Biomech* 2009, **42**(11):1622-1627.
71. Ruiz SA, Chen CS: **Emergence of Patterned Stem Cell Differentiation Within Multicellular Structures.** *Stem Cells* 2008, **26**(11):2921-2927.
72. Chen CS, Mrksich M, Huang S, Whitesides GM, Ingber DE: **Geometric control of cell life and death.** *Science* 1997, **276**(5317):1425-1428.
73. Juliano RL, Haskill S: **Signal transduction from the extracellular matrix.** *Journal of Cell Biology* 1993, **120**(3):577-585.
74. Maniotis AJ, Chen CS, Ingber DE: **Demonstration of mechanical connections between integrins, cytoskeletal filaments, and nucleoplasm that stabilize nuclear structure.** *Proc Natl Acad Sci USA* 1997, **94**(3):849-854.
75. Hynes RO: **Integrins: versatility, modulation, and signaling in cell adhesion.** *Cell* 1992, **69**(1):1-25.
76. Ingber DE: **Cellular tensegrity: defining new rules of biological design that govern the cytoskeleton.** *Journal of Cell Science* 1993, **104**(Pt 3):613-627.
77. Schmidt CE, Horwitz AF, Lauffenburger DA, Sheetz MP: **Integrin-cytoskeletal interactions in migrating fibroblasts are dynamic, asymmetric, and regulated.** *J Cell Biol* 1993, **123**(4):977-991.
78. Wang N, Butler JP, Ingber DE: **Mechanotransduction across the cell surface and through the cytoskeleton.** *Science* 1993, **260**(5111):1124-1127.
79. Urbich C, Dernbach E, Reissner A, Vasa M, Zeiher AM, Dimmeler S: **Shear stress-induced endothelial cell migration involves integrin signaling via the fibronectin receptor subunits alpha(5) and beta(1).** *Arteriosclerosis, Thrombosis & Vascular Biology* 2002, **22**(1):69-75.
80. Na S, Collin O, Chowdhury F, Tay B, Ouyang M, Wang Y, Wang N: **Rapid signal transduction in living cells is a unique feature of mechanotransduction.** *Proc Natl Acad Sci USA* 2008, **105**(18):6626-6631.
81. Clark CB, McKnight NL, Frangos JA: **Strain and strain rate activation of G proteins in human endothelial cells.** *Biochemical & Biophysical Research Communications* 2002, **299**(2):258-262.
82. Chachisvilis M, Zhang YL, Frangos JA: **G protein-coupled receptors sense fluid shear stress in endothelial cells.** *Proc Natl Acad Sci USA* 2006, **103**(42):15463-15468.
83. Janmey PA, Weitz DA: **Dealing with mechanics: mechanisms of force transduction in cells.** *Trends Biochem Sci* 2004, **29**(7):364-370.
84. Sigurdson W, Ruknudin A, Sachs F: **Calcium imaging of mechanically induced fluxes in tissue-cultured chick heart: role of stretch-activated ion channels.** *American Journal of Physiology* 1992, **262**(4 Pt 2):H1110-1115.
85. Munevar S, Wang YL, Dembo M: **Regulation of mechanical interactions between fibroblasts and the substratum by stretch-activated Ca²⁺ entry.** *J Cell Sci* 2004, **117**(Pt 1):85-92.
86. French AS, Stockbridge LL: **Potassium channels in human and avian fibroblasts.** *Proc R Soc Lond B Biol Sci* 1988, **232**(1269):395-412.
87. Wall ME, Banes AJ: **Early responses to mechanical load in tendon: role for calcium signaling, gap junctions and intercellular communication.** *J Musculoskelet Neuronal Interact* 2005, **5**(1):70-84.
88. Ruknudin A, Sachs F, Bustamante JO: **Stretch-activated ion channels in tissue-cultured chick heart.** *American Journal of Physiology* 1993, **264**(3 Pt 2):H960-972.
89. Malone AM, Anderson CT, Tummala P, Kwon RY, Johnston TR, Stearns T, Jacobs CR: **Primary cilia mediate mechanosensing in bone cells by a calcium-independent mechanism.** *Proc Natl Acad Sci USA* 2007, **104**(33):13325-13330.
90. Hinz B: **Masters and servants of the force: the role of matrix adhesions in myofibroblast force perception and transmission.** *Eur J Cell Biol* 2006, **85**(3-4):175-181.
91. Oberhauser AF, Badilla-Fernandez C, Carrion-Vazquez M, Fernandez JM: **The mechanical hierarchies of fibronectin observed with single-molecule AFM.** *J Mol Biol* 2002, **319**(2):433-447.
92. Friedland JC, Lee MH, Boettiger D: **Mechanically activated integrin switch controls alpha5beta1 function.** *Science* 2009, **323**(5914):642-644.

doi: 10.1186/1758-2555-2-16

Cite this article as: Wang and Li, Mechanics rules cell biology *Sports Medicine, Arthroscopy, Rehabilitation, Therapy & Technology* 2010, **2**:16

Submit your next manuscript to BioMed Central and take full advantage of:

- Convenient online submission
- Thorough peer review
- No space constraints or color figure charges
- Immediate publication on acceptance
- Inclusion in PubMed, CAS, Scopus and Google Scholar
- Research which is freely available for redistribution

Submit your manuscript at
www.biomedcentral.com/submit

